

CALL FOR PAPERS

DEADLINE: April 10, 2018

AQUA 2018 encourages the submission of high quality oral and poster presentations.

We strongly encourage authors to consider poster presentations because poster sessions will be an integral part of the program. Papers submitted for "oral presentation only" may not be accepted as oral presentations due to the limited number of available time slots. All abstracts must be in English-the official language of the conference.

Each oral presenter shall be entitled to no more than 20 minutes comprised of 15 minutes for a presentation, plus 5 minutes for questions. Authors of studies involving proprietary products or formulations should present this information in workshops or the trade show. Oral presentations can only use Power Point. Overhead projectors, slides and video players will not be available or allowed.

All presenters are required to pay their own registration, accommodation and travel expenses. AQUA 2018 cannot subsidize registration fees, travel or hotel costs.

No Abstract Book will be printed – a USB Abstract Book will be given to registered attendees.

INSTRUCTIONS FOR PREPARATION OF ABSTRACTS

Please refer to full guidelines at <https://www.was.org/EasOnline/abstracts/default.aspx>

- TITLE OF PAPER** : The paper title is printed in CAPITAL LETTERS, with the exception of scientific names which should be Upper/lower case and italicized. Scientific names should not be preceded or followed by commas or parentheses or other markings.
- AUTHOR(S)** : The first name should be the presenting author. Use *after the presenting author. Type in upper/lower case.
- ADDRESS AND EMAIL** : Type only the presenting author's institution, address and email. Type in upper/lower case.
- MAXIMUM LENGTH** : One Page
- PAGE SIZE** : Standard 210mm x 297mm A4 paper (portrait)
- MARGINS** : 2.5 cm margin throughout(left/right/top/bottom)
- SPACING** : Single spaced
- PARAGRAPHS** : Paragraphs should be separated by a blank line and should not be indented.
- FONT SIZES** : Character fonts should be 12 point type.
- FIGURES & TABLES** : Figures and tables are highly recommended. They should be reduced to the appropriate size for a one page abstract and should be clearly readable at the reduced size in black print only. The reduced figures and tables should be included in the abstract in camera-ready form.
- MEASUREMENTS** : Use metric units of measurement. When needed, English equivalents may be given in parentheses.

2.5 cm margin

EVALUATION OF JUVENILE AUSTRALIAN RED CLAW CRAYFISH *Decapoda quadricarinatus* FED PRACTICAL DIETS WITH AND WITHOUT SUPPLEMENTAL LECITHIN AND/OR CHOLESTEROL

Laura A. Mazinick*, Kenneth R. Thompson, Tracy Christian, Carl H. Webster, Lukman Manomczis, and David B. Rouse

Aquaculture Research Center
Kentucky State University
Frankfort, KY 40601
lmazinick@ksu.edu

Red claw crayfish (*Decapoda quadricarinatus*) are one of more than a hundred Australian freshwater crayfish species. However, because of its rapid growth rate, ease of spawning, wide temperature tolerance, and lack of a larval stage, red claw may be the best candidate for aquaculture. In the United States, red claw are only being investigated in an aquaculture species in the very early stages of development. Information exists on their nutritional requirements and practical diet formulations. No information exists on their requirements for lecithin and cholesterol to be added to their diet, these two nutrients are important for growth and cholesterol are very expensive. Since diet costs can be as much as 70% of the total cost of an aquaculture enterprise, it is imperative that the least expensive diet be formulated. The present study was conducted to determine if cholesterol and lecithin needs to be added to a practical diet for red claw crayfish.

An 8-week feeding trial was conducted in a recirculating system with newly-hatched juvenile (mean weight of 0.2 g) red claw, each stocked in six mesh culture units, individual units within fiberless tanks, each containing a water filter. Water was recirculated through mechanical filters. Water temperature was maintained at 27.20°C and lighting was provided by a 14-hour photoperiod. Diets were formulated to meet the minimum requirements for protein, lipid, carbohydrate, amino acids, vitamins, and minerals. Diets were fed three times per week. The goal of the study was to determine the effects of growth performance of newly hatched juvenile red claw when fed four practical diets with or without cholesterol and lecithin. Other practical diets included menhaden fish meal, soybean meal, shrimp meal, wheat flour, vitamin and mineral mix, pellet binder, cod liver oil and corn oil (Table 1).

After 8 weeks, red claw crayfish fed a practical diet without cholesterol had a significantly ($P < 0.05$) lower final weight, percentage weight gain, and specific growth rate (SGR) compared to crayfish fed all other diets (Table 2). These results indicate that a practical diet containing 2% cod liver oil and 1% corn oil and having no lecithin appears to be sufficient and that lecithin may not be necessary for juvenile red claw diets.

2.5 cm margin

21 cm long

2.5 cm margin

2.5 cm margin

21 cm wide

TABLE 1. Formulation of experimental diets fed to red claw crayfish.

	1	2	3	4
Menhaden Fish Meal	25.0	25.0	25.0	25.0
Soybean Meal	35.0	35.0	35.0	44.5
Lecithin 85	0.0	4.5	9.0	0.0
Cholesterol	1.0	1.0	0.0	0.0
Other	38.5	39.0	39.5	39.5

TABLE 2. Final weight, percentage weight gain, specific growth rate (SGR), and percentage survival of red claw crayfish fed four practical diets. Means in a column with different letters were significantly different ($P < 0.05$).

	1	2	3	4
Final weight (g)	4.97 ^a	8.00 ^a	3.60 ^b	5.11 ^a
Weight gain (%)	2344	2974	1713	2454
SGR (day ⁻¹)	5.76	5.66	4.68	5.41
Survival (%)	70	64	50	60

ALL ABSTRACTS AND PRESENTATIONS MUST BE IN ENGLISH

PLEASE SUBMIT YOUR ABSTRACT ONLINE

Submit abstracts via the internet at the meeting website : www.was.org

Follow the complete instructions on the online submission.

If you are unable to submit your abstract online, contact the Conference Manager for alternative methods at : worldaqua@was.org.